

A Simple Proof of the Classification of Normal Toeplitz Matrices

Akio Arimoto*

Abstract

We give an easy proof to show that every complex normal Toeplitz matrix is classified as either of type I or of type II. Instead of difference equations on elements in the matrix used in past studies, polynomial equations with coefficients of elements are used. In a similar fashion, we show that a real normal Toeplitz matrix must be one of four types: symmetric, skew-symmetric, circulant, or skew-circulant. Here we use trigonometric polynomials in the complex case and algebraic polynomials in the real case.

Key words. normal matrices, Toeplitz matrices. **AMS subject classifications.** 15A57, 47B15, 47B35

1 Introduction. Normal Toeplitz Matrices.

Ikramov[3] has firstly shown that a normal Toeplitz matrix over real field must be one of four types: symmetric, skew-symmetric (up to the principal diagonal), circulant, or skew-circulant. Later on the complex version of the problem has been solved by Ikramov[6] et.al and Gel'fgat[2]. The history of the theory has been described in [5]. The complex case was also proved by Farenick, Krupnik, Krupnik and Lee[1], and Ito[2] independently to [6]. We will give here another proof which, the author believes, is the simplest one. The complex case and the real case will be proved in a similar fashion, although trigonometric polynomials will be used in the complex case and algebraic polynomials will be used in the real case, respectively. Now denote

*Department of Electronics and Computer Engineering, Musashi Institute of Technology, Tamazutsumi 1-28-1, Setagaya-ku, Tokyo 158, Japan(arimoto@iname.com)

Toeplitz matrix of order $N + 1$ by T_N :

$$T_N = \begin{bmatrix} a_0 & a_{-1} & a_{-2} & \cdots & a_{-N} \\ a_1 & a_0 & a_{-1} & \cdots & a_{-(N-1)} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ a_{N-1} & a_{N-2} & a_{N-3} & \cdots & a_{-1} \\ a_N & a_{N-1} & a_{N-2} & \cdots & a_0 \end{bmatrix}.$$

T_N is called normal Toeplitz if $T_N T_N^* - T_N^* T_N = 0$, where T_N^* is a transposed conjugate of T_N . We can simply state classifications of normal Toeplitz matrices in the following way. A normal Toeplitz matrix T_N is of type I if and only if for some α_0 , with $|\alpha_0| = 1$,

$$(1) \quad \begin{bmatrix} a_{-1} & a_{-2} & \cdots & a_{-N} \end{bmatrix} = \alpha_0 \begin{bmatrix} \bar{a}_1 & \bar{a}_2 & \cdots & \bar{a}_N \end{bmatrix},$$

whereas T_N is of type II if and only if for some β_0 , with $|\beta_0| = 1$,

$$(2) \quad \begin{bmatrix} a_{-1} & a_{-2} & \cdots & a_{-N} \end{bmatrix} = \beta_0 \begin{bmatrix} a_N & a_{N-1} & \cdots & a_1 \end{bmatrix}.$$

Surely the complex case contains the real case, but when all the a'_k s are real valued, it will be found that α_0 and β_0 must be +1 or -1, hence we may have the further concrete classification of four types : A normal Toeplitz matrix T_N is symmetric if and only if

$$(3) \quad \begin{bmatrix} a_{-1} & a_{-2} & \cdots & a_{-N} \end{bmatrix} = \begin{bmatrix} a_1 & a_2 & \cdots & a_N \end{bmatrix},$$

T_N is skew-symmetric if and only if

$$(4) \quad \begin{bmatrix} a_{-1} & a_{-2} & \cdots & a_{-N} \end{bmatrix} = - \begin{bmatrix} a_1 & a_2 & \cdots & a_N \end{bmatrix},$$

T_N is circulant if and only if

$$(5) \quad \begin{bmatrix} a_{-1} & a_{-2} & \cdots & a_{-N} \end{bmatrix} = \begin{bmatrix} a_N & a_{N-1} & \cdots & a_1 \end{bmatrix},$$

T_N is skew-circulant if and only if

$$(6) \quad \begin{bmatrix} a_{-1} & a_{-2} & \cdots & a_{-N} \end{bmatrix} = - \begin{bmatrix} a_N & a_{N-1} & \cdots & a_1 \end{bmatrix}.$$

Throughout this paper, we assume a_0 to be zero without loss of generality since a_0 does not play any role in the above classifications.

2 Proof of Theorem in the Complex Case

Now we define trigonometric polynomials $s(x)$ and $t(x)$ with coefficients $a_1, a_2, \dots, a_N$ and $a_{-1}, a_{-2}, \dots, a_{-N}$, respectively:

$$s(x) = a_1 e^{ix} + a_2 e^{2ix} + \dots + a_N e^{Nix}$$

$$t(x) = a_{-1} e^{-ix} + a_{-2} e^{-2ix} + \dots + a_{-N} e^{-Nix}$$

Using these trigonometric polynomials, we can restate the conditions (1) and (2) in the previous section as

- (1') type I: $t(x) = \alpha_0 \overline{s(x)}$, with $|\alpha_0| = 1$,
(2') type II: $t(x) = \beta_0 s(x) e^{-i(N+1)x}$, with $|\beta_0| = 1$.

Theorem 1 *Every normal Toeplitz matrix is either of type I or of type II.*

Proof. Our proof is based on the expression in [7] p.998 or [1] p.1038. It is the necessary and sufficient conditions for T_N to be normal in terms of a_n that

$$(7) \quad a_m \bar{a}_n - \bar{a}_{-m} a_{-n} + \bar{a}_{N+1-m} a_{N+1-n} - a_{-(N+1-m)} \bar{a}_{-(N+1-n)} = 0,$$

$(1 \leq n, m \leq N).$

We now rewrite (7) by using polynomials $s(x)$ and $t(x)$. Multiply both sides of (7) by $e^{imx} e^{-iny}$ and sum these new expressions over all m and n to obtain

$$(8) \quad s(x) \overline{s(y)} - \overline{t(x)} t(y) + \overline{s(x)} s(y) e^{i(N+1)(x-y)} - t(x) \overline{t(y)} e^{i(N+1)(x-y)} = 0.$$

It easily seen that (8) is equivalent to (7), and (1') or (2') implies (8). Hence both (1') and (2') are sufficient conditions for T_N to be normal. So it only remains to show the necessity of the condition. When we take $x = y$ in (8), we have

$$(9) \quad s(x) \overline{s(x)} - t(x) \overline{t(x)} = 0, \text{ or } |s(x)| = |t(x)|$$

for each x . Hence except for trivial case $t(x) \equiv 0$, we can find an x_0 such that $t(x_0) \neq 0$ and there exist α and β such that $s(x_0) = \alpha t(x_0)$ and $t(x_0) = \beta \overline{t(x_0)}$, with $|\alpha| = |\beta| = 1$. By setting $y = x_0$ and then by dividing through by $t(x_0)$, equation (8) becomes

$$(10) \quad s(x) \bar{\alpha} - \overline{t(x)} \beta + \overline{s(x)} \alpha \beta e^{i(N+1)(x-x_0)} - t(x) e^{i(N+1)(x-x_0)} = 0$$

from which we have

$$(11) \quad \overline{t(x)} = s(x) \overline{\alpha\beta} + \overline{s(x)} \alpha e^{i(N+1)(x-x_0)} - t(x) \bar{\beta} e^{i(N+1)(x-x_0)}.$$

Substituting the right hand side of (11) into $\overline{t(x)}$ of (9), we have

$$(12) \quad \left\{ s(x) - t(x) \alpha e^{i(N+1)(x-x_0)} \right\} \left\{ \overline{s(x)} - t(x) \overline{\alpha\beta} \right\} = 0,$$

which implies our required results if we take $\alpha_0 = \alpha\beta$ and $\beta_0 = \bar{\alpha} e^{i(N+1)x_0}$. We should notice here that (12) is the product of polynomials. If one of the polynomials in (12) is nonzero at $x = x_0$, then by continuity the polynomial is nonzero in an open neighbourhood $\mathcal{U}$ of x_0 , which implies that the other polynomial in the product (12) is identically zero on $\mathcal{U}$. Because the only polynomial with a continuum of roots is the zero polynomial, equation (12) implies equation (1') or (2'). Hence type I or type II must occur.

3 Proof of Theorem in the Real Case.

Theorem 2 *Every real normal Toeplitz matrix is either symmetric, skew-symmetric, circulant or skew-circulant.*

Proof. We define algebraic polynomials $p(x)$ and $q(x)$ such that

$$p(x) = a_1x + a_2x^2 + \cdots + a_Nx^N$$

$$q(x) = a_{-1}x + a_{-2}x^2 + \cdots + a_{-N}x^N$$

Also in this real case we can apply (7) in the previous section as necessary and sufficient conditions for T_N to be normal in terms of real valued a_n :

$$(13) \quad a_m a_n - a_{-m} a_{-n} + a_{N+1-m} a_{N+1-n} - a_{-(N+1-m)} a_{-(N+1-n)} = 0 \quad (1 \leq n, m \leq N)$$

Now we introduce the reciprocal polynomials to $p(x)$ and $q(x)$

$$\tilde{p}(x) = a_N x + a_{N-1} x^2 + \cdots + a_1 x^N$$

and

$$\tilde{q}(x) = a_{-N} x + a_{-N+1} x^2 + \cdots + a_{-1} x^N,$$

respectively. Multiply both sides of equation (13) by $x^m y^n$ and then sum the resulting expressions over all m and n to obtain the following analogue of (8):

$$(14) \quad p(x) p(y) - q(x) q(y) + \tilde{p}(x) \tilde{p}(y) - \tilde{q}(x) \tilde{q}(y) = 0.$$

Noticing the relation $\tilde{q}(x) = x^{N+1}q(1/x)$ and $\tilde{p}(x) = x^{N+1}p(1/x)$, and letting $y = 1/x$ in (14), we have $p(x)\tilde{p}(x) = q(x)\tilde{q}(x)$. By setting $x = y$ and then multiplying both sides of (14) by $p^2(x)$, we have from the relation $p^2(x)\tilde{p}^2(x) = q^2(x)\tilde{q}^2(x)$,

$$(15) \quad p^4(x) - p^2(x)q^2(x) + q^2(x)\tilde{q}^2(x) - p^2(x)\tilde{q}^2(x) = 0,$$

which can be factorized as the following way:

$$(16) \quad (p^2(x) - q^2(x))(p^2(x) - \tilde{q}^2(x)) = 0.$$

From this formula, we obtain four types, symmetric $p(x) = q(x)$, skew-symmetric $p(x) = -q(x)$, circulant $p(x) = \tilde{q}(x)$ and skew-circulant $p(x) = -\tilde{q}(x)$, which are equivalent to (3),(4),(5) and (6), respectively.

References

- [1] D.R.Farenick, M.Krupnik,N.Krupnik, and W.Y.Lee, Normal Toeplitz matrices, SIAM J. Matrix Anal. Appl.,17(1996),pp.1037-1043
- [2] V.I.Gel'FGAT, A normality criterium for Toeplitz matrices, Zh.Vychisl.Mat.i Fiz.,35(1995)pp.1428-1432(in Russian). Comput.Math.Phys.,35(1995),pp.1147-1150
- [3] Kh.D.Ikramov, Describing normal Toeplitz matrices, Zh.Vychisl.Mat.i Fiz.,34(1994)pp.473-479 (in Russian). Comput.Math.Phys.,34(1994),pp.399-404
- [4] Kh.D.Ikramov, Classification of normal Toeplitz matrices with real entries, Zh.Vychisl.Mat.iFiz.,57(1995)pp.1428-1432(in Russian). Comput.Math.Phys.,34(1994),pp.399-404
- [5] Kh.D.Ikramov, Comments on Normal Toeplitz Matrices by Farenick et al., SIAM J.Matrix Anal.Appl.,vol.18(1997),no.2,p.518
- [6] Kh.D.Ikramov and V.N.Chugunov, Normality condition for a complex Toeplitz matrix, Zh.Vychisl.Mat.i.Fiz.,36(1996)pp.3-10(in Russian)Comput.Math.Phys.,34(1996),pp.131-137
- [7] T.Ito, Every normal Toeplitz matrix is either of type I or of type II, SIAM J.Matrix Anal.Appl.,17(1996),pp.998-1006